



Polynomial Regression

$$y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \beta_3 x_i^3 + \dots + \beta_d x_i^d + \epsilon_i$$

Degree-4 Polynomial



Details

- Create new variables $X_1 = X$, $X_2 = X^2$, etc and then treat as multiple linear regression.
- Not really interested in the coefficients; more interested in the fitted function values at any value x_0 :

$$\hat{f}(x_0) = \hat{\beta}_0 + \hat{\beta}_1 x_0 + \hat{\beta}_2 x_0^2 + \hat{\beta}_3 x_0^3 + \hat{\beta}_4 x_0^4.$$

- Since $\hat{f}(x_0)$ is a linear function of the $\hat{\beta}_\ell$, can get a simple expression for *pointwise-variances* $\text{Var}[\hat{f}(x_0)]$ at any value x_0 . In the figure we have computed the fit and pointwise standard errors on a grid of values for x_0 . We show $\hat{f}(x_0) \pm 2 \cdot \text{se}[\hat{f}(x_0)]$.
- We either fix the degree d at some reasonably low value, else use cross-validation to choose d .

Details continued

- Logistic regression follows naturally. For example, in figure we model

$$\Pr(y_i > 250|x_i) = \frac{\exp(\beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \dots + \beta_d x_i^d)}{1 + \exp(\beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \dots + \beta_d x_i^d)}.$$



- To get confidence intervals, compute upper and lower bounds on *on the logit scale*, and then invert to get on probability scale.
- Can do separately on several variables—just stack the variables into one matrix, and separate out the pieces afterwards (see GAMs later).
- Caveat: polynomials have notorious tail behavior — very bad for extrapolation.
- Can fit using $\mathbf{y} \sim \text{poly}(\mathbf{x}, \text{degree} = 3)$ in formula.

Step Functions

Another way of creating transformations of a variable — cut the variable into distinct regions.

$$C_1(X) = I(X < 35), \quad C_2(X) = I(35 \leq X < 65), \dots, C_3(X) = I(X \geq 65)$$

Piecewise Constant



Step functions continued

- Easy to work with. Creates a series of dummy variables representing each group.
- Useful way of creating interactions that are easy to interpret. For example, interaction effect of **Year** and **Age**:

$$I(\text{Year} < 2005) \cdot \text{Age}, \quad I(\text{Year} \geq 2005) \cdot \text{Age}$$

would allow for different linear functions in each age category.

- In R: `I(year < 2005)` or `cut(age, c(18, 25, 40, 65, 90))`.
- Choice of cutpoints or *knots* can be problematic. For creating nonlinearities, smoother alternatives such as *splines* are available.


Piecewise Polynomials

- Instead of a single polynomial in X over its whole domain, we can rather use different polynomials in regions defined by knots. E.g. (see figure)


$$y_i = \begin{cases} \beta_{01} + \beta_{11}x_i + \beta_{21}x_i^2 + \beta_{31}x_i^3 + \epsilon_i & \text{if } x_i < c; \\ \beta_{02} + \beta_{12}x_i + \beta_{22}x_i^2 + \beta_{32}x_i^3 + \epsilon_i & \text{if } x_i \geq c. \end{cases}$$

- Better to add constraints to the polynomials, e.g. continuity.
- *Splines* have the “maximum” amount of continuity.


Piecewise Cubic




Continuous Piecewise Cubic



Cubic Spline



Linear Spline



Linear Splines

A linear spline with knots at ξ_k , $k = 1, \dots, K$ is a piecewise linear polynomial continuous at each knot.

We can represent this model as



$$y_i = \beta_0 + \beta_1 b_1(x_i) + \beta_2 b_2(x_i) + \dots + \beta_{K+3} b_{K+3}(x_i) + \epsilon_i,$$

where the b_k are *basis functions*.

$$\begin{aligned} b_1(x_i) &= x_i \\ b_{k+1}(x_i) &= (x_i - \xi_k)_+, \quad k = 1, \dots, K \end{aligned}$$

Here the $(\)_+$ means *positive part*; i.e.

$$(x_i - \xi_k)_+ = \begin{cases} x_i - \xi_k & \text{if } x_i > \xi_k \\ 0 & \text{otherwise} \end{cases}$$



Cubic Splines

A cubic spline with knots at ξ_k , $k = 1, \dots, K$ is a piecewise cubic polynomial with continuous derivatives up to order 2 at each knot.

Again we can represent this model with truncated power basis functions

$$y_i = \beta_0 + \beta_1 b_1(x_i) + \beta_2 b_2(x_i) + \dots + \beta_{K+3} b_{K+3}(x_i) + \epsilon_i,$$

$$b_1(x_i) = x_i$$

$$b_2(x_i) = x_i^2$$

$$b_3(x_i) = x_i^3$$

$$b_{k+3}(x_i) = (x_i - \xi_k)_+^3, \quad k = 1, \dots, K$$

where

$$(x_i - \xi_k)_+^3 = \begin{cases} (x_i - \xi_k)^3 & \text{if } x_i > \xi_k \\ 0 & \text{otherwise} \end{cases}$$

